

LIGHT HADRON MASSES ON COARSE LATTICES WITH IMPROVED ACTIONS

FRANK X. LEE,^a DEREK B. LEINWEBER^b

^a*TRIUMF, 4004 Wesbrook Mall, Vancouver, British Columbia, Canada V6T 2A3*

^b*Department of Physics, University of Washington, Seattle, WA 98195*

The advent of mean-field improved perturbation theory¹ has generated renewed interest in the on-shell improvement of lattice QCD actions. The remarkable success of the $\mathcal{O}(a^2)$ -improved gauge-field action² has led us to reconsider the two-link $\mathcal{O}(a^2)$ -improved fermion action of Hamber and Wu,³ with mean-field improved operators.

Tree-level $\mathcal{O}(a^2)$ -improved mean-field improved gauge configurations are generated in the quenched approximation via the Cabibbo-Marinari pseudo-heat-bath algorithm. Fermion propagators are obtained via the Stabilized Biconjugate Gradient algorithm.⁴ We consider a local source and both local and gauge-invariant Gaussian smeared sinks.⁵ We found smearing is necessary in order to extract observables prior to a loss of signal. Five quark masses ranging from the strange quark mass to one-third the strange quark mass are considered. The strange quark mass was taken to be approximately 180 MeV.

To evaluate scaling in the observables, two lattice spacings a are considered on equal physical volume lattices. The lattice parameters are summarized in table 1. A single-elimination jackknife estimate of the covariance matrix provides a correlated χ^2 which is used to select the time-slice analysis regime.

Dispersion is examined by calculating the speed of light squared from the relation $c^2 = (E_p^2 - M^2)/p^2$, to be compared with 1. E_p and M are extracted from lattice correlation functions and p is the smallest nontrivial momentum $(1, 0, 0) 2\pi/L$. Table 2 compares ρ -meson results for $m_q \sim 125$ MeV with the $\mathcal{O}(a)$ -improved SW action⁶ and the $\mathcal{O}(a^2)$ -improved D234 action⁷. The last column tests rotational symmetry for the Hamber Wu action (HW) by determining c^2 from the smallest two-dimensional diagonal momentum. The $\mathcal{O}(a^2)$ -improved actions perform well at the spacing of 0.24 fm.

Hadron mass ratios within the baryon octet, decuplet and the vector meson nonet scale well and look very similar to experiment. However, mass ratios of hadrons with differing spins reveal less than satisfactory scaling for the

Table 1: Summary of lattice parameters.

Volume	β	a_{st}	N_U	u_0	κ_{cr}
$6^3 \times 12$	6.25	0.40 fm	90	0.820	0.1967(2)
$10^3 \times 16$	7.00	0.24 fm	37	0.866	0.1823(3)

Table 2: Evaluation of dispersion and rotational invariance via c^2 .

a_{st}	SW (1,0,0)	D234 (1,0,0)	HW (1,0,0)	HW (1,1,0)
0.40	0.48(4)	0.93(3)	0.88(6)	0.81(9)
0.24	—	1.00(6)	0.96(9)	1.03(7)

larger lattice spacing. The N/ρ mass ratio summarized in figure 1 is the most revealing ratio. The Hamber and Wu action has reproduced the state of the art quenched QCD ratio on a tiny $10^3 \times 16$ lattice. This result bodes well for future explorations of hadron phenomenology beyond the quenched approximation.




Figure 1: The N/ρ mass ratio as a function of the lattice spacing a_ρ for a variety of lattice QCD actions. Solid symbols denote the standard Wilson action. Open symbols denote improved actions including SW⁶ ($\diamond$), D234⁷ ($\triangle$), HW⁸ ($\square$), and HW ($\circ$) (this work).

This work is supported in part by the Natural Sciences and Engineering Research Council of Canada and U.S. DOE under grant DE-FG06-88ER40427.

1. G. P. Lepage and P. B. Mackenzie, Phys. Rev. D **48**, 2250 (1993).
2. M. Alford *et al.*, Phys. Lett. **B361**, 87 (1995).
3. W. Hamber and C. Wu, Phys. Lett. **133B**, 351 (1983).
4. A. Frommer *et al.*, J. Mod. Phys. **C5**, 1073 (1994).
5. S. Güsken, Nucl. Phys. B (Proc. Suppl.) **17**, 361 (1990).
6. S. Collins, *et al.*, Nucl. Phys. B (Proc. Suppl.) **47**, 366 (1995).
7. M. Alford, *et al.*, Nucl. Phys. B (Proc. Suppl.) **47**, 370 (1995).
8. H. R. Fiebig and R. M. Woloshyn, TRI-PP-96-1, hep-lat/9603001.